

DESOAGEN[®] WT-H

Major component

- Anionic Surfactant

Property

- Rapid and even wetting, and remove dirt and fats when used for soaking.
- Promote penetration of chemicals, uniform swelling of pelt and give clean grain, when used for liming.
- Emulsify and disperse natural fats effectively when used in deliming and bating.
- Fast wetting for conditioning of wet-blue or crust.

Specification

Appearance(25°C):	Viscous liquid
Color:	Yellowish
pH(25°C):	9.5±1.0
Ionicity:	Anionic
APEO:	Comply with EU REACH laws and regulations

Application

- Good compatibility with other chemicals. Could be added together with other chemicals.
- Soaking (in terms of wet salted): **DESOAGEN WT-H** 0.3~0.5%
- Liming (in terms of wet salted): **DESOAGEN WT-H** 0.4~0.5%
- Deliming and Bating (in terms of limed pelt):**DESOAGEN WT-H** 0.3~0.5%
- Conditioning of wet blue or crust (in terms of shaved wet-blue): **DESOAGEN WT-H** 0.5~1.0%

DESOAGEN[®] WT-H

Safety

- This is not a hazardous chemical.
- Observe all the precautions required for handling chemicals.
- Read attentively and observe all the recommendations in our **Material Safety Data Sheet**.

Transportation and Storage

- The product has a shelf life of at least two years if it is stored in its tightly sealed original packaging at temperatures between 5°C and 35°C.
- This product should be stored in a cool and dry place, lidded and keep upright. Avoid direct exposure to the sun or frost.
- Viscosity changes along with temperature. Higher temperature leads to lower viscosity.
- Separation or precipitation might occur when temperature is low. Stir before use and properties will not be compromised.
- Re-seal the package immediately after using, use up as soon as possible once the package is opened.
- Read attentively and observe all the recommendations in our **Material Safety Data Sheet**.